

Landenoverzicht exporteisen Zaaizaden.

Land: **Israël**

Overzicht van de laatste wijziging(en)

versie	datum	toelichting
3.88	01-04-2026	Vervanging oude code <i>Lathyrus odoratus</i>
3.87	25-03-2026	Toevoeging nieuwe code <i>Petroselinum sp., Phaseolus sp.</i> , correctie codes m.b.t. 'test on-arrival'
3.86	28-01-2026	Correctie peststatus ToMV

Landenoverzicht exporteisen Zaaizaden - Israël

ISRAËL (IL)	Certificaat Export	Certificaat re-export	Taal	Grondeis	Invoervergunning
	1	20	E	-	meestal

Algemene informatie

Geleidelijk gaan we over naar een nieuwe coderingssysteem waardoor, voorlopig, twee verschillende systemen in gebruik zijn in dit document.

Uitleg codes, zie NVWA-site: <https://www.nvwa.nl/onderwerpen/export-planten-groenten-fruit-plantaardige-producten/documenten/export/fytosanitair/voorschriften/algemeen/toelichting-landeneisen>

Legenda

\$ = zie Register Dekkingen Zaaizaden

~ = de dekking van dit organisme moet nog bepaald worden. De datasheet moet nog worden opgesteld. Houd aub er rekening mee dat dit een paar weken in beslag kan nemen

* = zie 'Invoerverbod'

Pre-exportcertificaten

De exporteur dient er rekening mee te houden dat sommige garanties moeten worden afgegeven door het EU-land van origine. Dit gebeurt met een pre-exportcertificaat. Op dit certificaat verklaart het EU land van origine dat de planten, plantaardige producten of andere materialen die geteeld, geproduceerd, opgeslagen of verwerkt zijn, voldoen aan specifieke fytosanitaire voorschriften met betrekking tot één of meer van de volgende aspecten:

- de afwezigheid of aanwezigheid van bepaalde organismen;
- de oorsprong in of op een specifiek veld, productiefaciliteit, productieplaats of gebied;
- de status van een plaagorganisme in het veld, in de productiefaciliteit, op de productieplaats, in het gebied of het land van oorsprong;
- het resultaat van inspecties, het nemen van monsters van en het uitvoeren van tests.

Zie ook:

<https://www.nvwa.nl/documenten/export/fytosanitair/voorschriften/toelichting-certificaten/instructie-waarmeden-fytosanitair-certificaat>

Algemene eisen

Bron

Plant Import Regulations, February 2009 (Ministry of Agriculture and Rural Development, Plant Protection and Inspection)

Treatment Manual:

<https://www.gov.il/BlobFolder/reports/treatment->

Landenoverzicht exporteisen Zaaizaden.

Land: **Israël**

[manual/he/import import plants and plant products treatment-manual-version-2%2011.05.2016.pdf](#)

Certificaat vereist voor:

Alle gewassoorten.

Invoerverbod

Gewassoorten aangemerkt met een asterisk (*) in het document met de fyto-sanitaire wet- en regelgeving van Israël zijn uitsluitend toegestaan indien uit de volgende origines: Australië, België, Canada, Denemarken, Duitsland, Finland, Frankrijk, Ierland, Israël, Italië, Japan, Nederland, Nieuw-Zeeland, Oostenrijk, Portugal, Spanje, Taiwan, Verenigd Koninkrijk, Verenigde Staten, Zweden, Zwitserland.

Soms wordt er d.m.v. invoervergunningen bekend gemaakt dat andere origines ook zijn toegestaan. Deze worden dan ook genoemd bij de origines van het product, onder het kopje 'Bijschrijvingen'.

Verder, voor zover bekend, geen invoerverbod voor zaaizaden.

Certificeringseisen

Capsicum annum, Solanum lycopersicum en Solanum melongena:

Bij re-export moet de garantie van het origineel FC worden overgenomen. Indien er géén garantie is afgegeven moet er een labtoets plaatsvinden. Deze toets moet op het re-export FC worden vermeld in de bijschrijving ("The seeds have been tested in an official laboratory and found free from....").

De zaden van bovengenoemde gewassen worden na aankomst in Israël getoetst op de bij de producten genoemde organismen in een officieel laboratorium.

Bijschrijvingen

Legenda:

- a1 = The seeds have been tested in an official laboratory and found free from
- a2 = The seeds will be tested on-arrival to proof freedom of
- b1 = The seeds originate from plants that were inspected during the active growing season and found free from
- b2 = The seeds originate from a field inspected during the active growing season and found free from
- b3 = The place of production was officially inspected during the growing season and found free from
- c1 = The seeds are from an area known to be free from
- d1 = The country of origin is free from
- d2 = The seeds originate from plants grown in a country known to be free from
- e1 = The seeds originate from plants that have been tested in an official laboratory and found free from

Allium sp. *

Origines Argentinië, Australië, België, Canada, Denemarken, Duitsland, Finland, Frankrijk, Ierland, Israël, Italië, Japan, Nederland, Nieuw-Zeeland, Oostenrijk, Portugal, Spanje, Taiwan, Verenigd Koninkrijk, Verenigde Staten, Zweden, Zwitserland

(a1) The seeds have been tested in an official laboratory and found free from *Ditylenchus dipsaci* (2), *Sclerotium cepivorum* (21), *Urocystis cepulae* (2).

OF

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Ditylenchus dipsaci* (6), *Sclerotium cepivorum* (6), *Urocystis cepulae* (6).

OF

Landenoverzicht exporteisen Zaaizaden.

Land: **Israël**

(a2) The seeds will be tested on-arrival to proof freedom of *Ditylenchus dipsaci*, *Sclerotium cepivorum*, *Urocystis cepulae*.

Origine Zuid-Afrika (permits)

(d1) The country of production is free from: *Urocystis cepulae* (4-NL niet vrij).

EN

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Ditylenchus dipsaci* (6), *Stromatinia cepivora* (6), *Xanthomonas axonopodis* pv. *allii* (6), *Pantoea ananatis* (6) and tobacco rattle virus (6).

OF

(a1) The seeds have been tested in an official laboratory and found free from *Ditylenchus dipsaci* (2), *Stromatinia cepivora* (21), *Xanthomonas axonopodis* pv. *allii* (2), *Pantoea ananatis* (2) and tobacco rattle virus (2).

Anethum graveolens *

Origines Australië, België, Canada, Denemarken, Duitsland, Finland, Frankrijk, Ierland, Israël, Italië, Japan, Nederland, Nieuw-Zeeland, Oostenrijk, Portugal, Spanje, Taiwan, Verenigd Koninkrijk, Verenigde Staten, Zweden, Zwitserland

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Phymatotrichopsis ominivora* (IV of I), artichoke yellow ringspot virus (6).

OF

(a1) The seeds have been tested in an official laboratory and found free from *Phymatotrichopsis ominivora* (IV of I), artichoke yellow ringspot virus (2).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Phymatotrichopsis ominivora*, artichoke yellow ringspot virus.

Antirrhinum sp.

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Puccinia antirrhini* (I), *Pseudomonas syringae* pv. *antirrhini* (6).

OF

(a1) The seeds have been tested in an official laboratory and found free from *Puccinia antirrhini* (I), *Pseudomonas syringae* pv. *antirrhini* (2).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Puccinia antirrhini*, *Pseudomonas syringae* pv. *antirrhini*.

Apium graveolens *

Origines Australië, België, Canada, Denemarken, Duitsland, Finland, Frankrijk, Ierland, Israël, Italië, Japan, Nederland, Nieuw-Zeeland, Oostenrijk, Portugal, Spanje, Taiwan, Verenigd Koninkrijk, Verenigde Staten, Zweden, Zwitserland

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Phyllosticta apii* (I of IV), *Phytophthora cryptogea* (I of IV), *Pseudomonas syringae* pv. *apii* (6), strawberry latent ringspot virus (6).

OF

(a1) The seeds have been tested in an official laboratory and found free from *Phyllosticta apii* (I of IV), *Phytophthora cryptogea* (I of IV), *Pseudomonas syringae* pv. *apii* (2), strawberry latent ringspot virus (2).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Phyllosticta apii*, *Phytophthora cryptogea*, *Pseudomonas syringae* pv. *apii*, strawberry latent ringspot virus.

Landenoverzicht exporteisen Zaaizaden.

Land: **Israël**

EN

Seeds are free from noxious weed seeds of *Cirsium arvense* (I).

Asparagus sp.

The seeds are free of pulp.

Atriplex sp. *

Origines Australië, België, Canada, Denemarken, Duitsland, Finland, Frankrijk, Ierland, Israël, Italië, Japan, Nederland, Nieuw-Zeeland, Oostenrijk, Portugal, Spanje, Taiwan, Verenigd Koninkrijk, Verenigde Staten, Zweden, Zwitserland

(b1) + *Microsphaeropsis olivaceae* (~).

Bellis sp.

(a1) The seeds have been tested in an official laboratory and found free from *Phoma bellidis* (~)

OF

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Phoma bellidis* (~).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Phoma bellidis*.

Beta vulgaris *

Origines Australië, België, Canada, Denemarken, Duitsland, Finland, Frankrijk, Ierland, Israël, Italië, Japan, Nederland, Nieuw-Zeeland, Oostenrijk, Portugal, Spanje, Taiwan, Verenigd Koninkrijk, Verenigde Staten, Zweden, Zwitserland

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Ditylenchus dipsaci* 6), *Curtobacterium flaccumfaciens* pv. *betae* (6), beet cryptic virus 1 (6), beet cryptic virus 2 (6).

Brassica juncea *, **Brassica napus ***, **Brassica oleracea ***, **Brassica pekinensis ***, **Brassica rapa *** + Chili

Origines Australië, België, Canada, Chili, Denemarken, Duitsland, Finland, Frankrijk, Ierland, Israël, Italië, Japan, Nederland, Nieuw-Zeeland, Oostenrijk, Portugal, Spanje, Taiwan, Verenigd Koninkrijk, Verenigde Staten, Zweden, Zwitserland

(a1) The seeds have been tested in an official laboratory and found free from *Leptosphaeria maculans* (2), *Xanthomonas campestris* pv. *campestris* (2).

OF

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Leptosphaeria maculans* (6), *Xanthomonas campestris* pv. *campestris* (6).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Leptosphaeria maculans*, *Xanthomonas campestris* pv. *campestris*.

Cactaceae

(a1 of a2) + *Helminthosporium cactivorum* (I) and seeds are free from pulp.

Calendula officinalis

(a1) The seeds have been tested in an official laboratory and found free from *Drechlera hawaiiensis* (2).

OF

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Drechlera hawaiiensis* (6).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Drechlera hawaiiensis*.

Landenoverzicht exporteisen Zaaizaden.

Land: **Israël**

Callistephus chinensis

(a1) The seeds have been tested in an official laboratory and found free from *Aphelenchoides ritzemabosi* (I), *Fusarium oxysporum* f. sp. *callistephi* (2), tomato aspermy virus (2).

OF

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Aphelenchoides ritzemabosi* (IV), *Fusarium oxysporum* f. sp. *callistephi* (6), tomato aspermy virus (6).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Aphelenchoides ritzemabosi*, *Fusarium oxysporum* f. sp. *callistephi*, tomato aspermy virus.

Capsicum annuum

Zie ook 'Certificeringseisen'.

Eis zonder bijschrijving:

Partij vrij van tomato brown rugose fruit virus (2).

EN

The production area is free from *Phytophthora capsici* (4-NL niet vrij; 5), *Clavibacter michiganensis* subsp. *michiganensis* (4-NL vrij; 5), *Erwinia carotovora* subsp. *carotovora* (4-NL niet vrij; 5), *Pseudomonas corrugata* (4-NL niet vrij; 5), *Xanthomonas vesicatoria* (4-NL vrij; 5), tobacco mosaic virus (4-NL niet vrij; 5), pepper mild mottle virus (4-NL niet vrij; 5), tobacco mild green mosaic virus (4-NL vrij; 5), tomato black ring virus (4-NL niet vrij; 5), tomato ringspot virus (4-NL niet vrij; 5).

OF

The consignment was checked during active growth and it was found free from *Phytophthora capsici* (6), *Clavibacter michiganensis* subsp. *michiganensis* (6), *Erwinia carotovora* subsp. *carotovora* (6), *Pseudomonas corrugata* (6), *Xanthomonas vesicatoria* (6), tobacco mosaic virus (6), pepper mild mottle virus (6), tobacco mild green mosaic virus (6), tomato black ring virus (6), tomato ringspot virus (6).

OF

Based on laboratory tests the consignment was found free from *Phytophthora capsici* (2), *Clavibacter michiganensis* subsp. *michiganensis* (2), *Erwinia carotovora* subsp. *carotovora* (2), *Pseudomonas corrugata* (2), *Xanthomonas vesicatoria* (2), tobacco mosaic virus (2), pepper mild mottle virus (2), tobacco mild green mosaic virus (2), tomato black ring virus (2), tomato ringspot virus (2). (*Let op: labuitslag meesturen*).

Carthamus tinctorius *

Origines Australië, België, Canada, Denemarken, Duitsland, Finland, Frankrijk, Ierland, Israël, Italië, Japan, Nederland, Nieuw-Zeeland, Oostenrijk, Portugal, Spanje, Taiwan, Verenigd Koninkrijk, Verenigde Staten, Zweden, Zwitserland

(a1) The seeds have been tested in an official laboratory and found free from *Cercospora carthami* (I), *Fusarium oxysporum* f. sp. *carthami* (2), *Puccinia carthami* (I), *Verticillium albo-atrum* (I).

OF

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Cercospora carthami* (IV), *Fusarium oxysporum* f. sp. *carthami* (6), *Puccinia carthami* (I), *Verticillium albo-atrum* (IV).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Cercospora carthami*, *Fusarium oxysporum* f. sp. *carthami*, *Puccinia carthami*, *Verticillium albo-atrum*.

Centaurea cyanus

(a1) The seeds have been tested in an official laboratory and found free from *Phoma* spp (I).

OF

Landenoverzicht exporteisen Zaaizaden.

Land: **Israël**

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Phoma* spp. (IV)

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Phoma* spp.

Chenopodium sp.

(e1) The seeds originate from plants that have been tested in an official laboratory and found free from *Ascochyta hyalospora* (9), *Hibiscus latent ringspot virus* (9), *Raspberry bushy dwarf virus* (9), *Sowbane mosaic virus* (9).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Ascochyta hyalospora*, *Hibiscus latent ringspot virus*, *Raspberry bushy dwarf virus*, *Sowbane mosaic virus*.

Chrysanthemum sp.

(a1 of a2 of b) + *Aphelenchoides ritzemabosi* (I), *Didymella ligulicola* (I of IV), *Peronospora radii* (I of IV), *Erwinia chrysanthemi* (I).

(a1) The seeds have been tested in an official laboratory and found free from *Aphelenchoides ritzemabosi* (I), *Didymella ligulicola* (I of IV), *Peronospora radii* (I of IV), *Erwinia chrysanthemi* (2).

OF

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Aphelenchoides ritzemabosi* (IV), *Didymella ligulicola* (IV), *Peronospora radii* (IV), *Erwinia chrysanthemi* (6).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Aphelenchoides ritzemabosi*, *Didymella ligulicola*, *Peronospora radii*, *Erwinia chrysanthemi*.

Cicer sp. *

Origines Australië, België, Canada, Denemarken, Duitsland, Finland, Frankrijk, Ierland, Israël, Italië, Japan, Nederland, Nieuw-Zeeland, Oostenrijk, Portugal, Spanje, Taiwan, Verenigd Koninkrijk, Verenigde Staten, Zweden, Zwitserland

(b2 of a1 of a2) + *Ascochyta rabiei* (~), *Fusarium oxysporum* f. sp. *ciceris* (2; 6), broad bean mottle virus (~), pea mosaic virus = bean yellow mosaic virus (~).

Cichorium sp. *

Origines Australië, België, Canada, Denemarken, Duitsland, Finland, Frankrijk, Ierland, Israël, Italië, Japan, Nederland, Nieuw-Zeeland, Oostenrijk, Portugal, Spanje, Taiwan, Verenigd Koninkrijk, Verenigde Staten, Zweden, Zwitserland

(a1) The seeds have been tested in an official laboratory and found free from *Alternaria cichorii* (I), *Fusarium avenaceum* (I) and *Gibberella avenacea* (I).

OF

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Alternaria cichorii* (IV), *Fusarium avenaceum* (IV) and *Gibberella avenacea* (IV)

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Alternaria cichorii*, *Fusarium avenaceum* and *Gibberella avenacea*.

Citrullus lanatus * + Chili, Peru

Origines Australië, België, Canada, Denemarken, Duitsland, Finland, Frankrijk, Ierland, Israël, Italië, Japan, Nederland, Nieuw-Zeeland, Oostenrijk, Portugal, Spanje, Taiwan, Thailand, Verenigd Koninkrijk, Verenigde Staten, Zweden, Zwitserland

(c1) The seeds are from an area known to be free from *Acidovorax avenae* pv *citrulli* (4 NL vrij; 5).

EN

Landenoverzicht exporteisen Zaaizaden.

Land: **Israël**

(a1) The seeds have been tested in an official laboratory and found free from *Acidovorax avenae* pv *citrulli* (2)

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Acidovorax avenae* pv *citrulli*.

EN

(b1) The seeds originate from plants that were inspected during active growing season and found free from cucumber green mottle mosaic virus (6), squash mosaic virus (6), *Didymella bryoniae* (6), *Fusarium oxysporum* f. sp. *niveum* (6), *Colletotrichum orbiculare* (6).

OF

(a1) The seeds have been tested in an official laboratory and found free from cucumber green mottle mosaic virus (2), squash mosaic virus (2), *Didymella bryoniae* (2), *Fusarium oxysporum* f. sp. *niveum* (2), *Colletotrichum orbiculare* (2).

OF

(a2) The seeds will be tested on-arrival to proof freedom of cucumber green mottle mosaic virus, squash mosaic virus, *Didymella bryoniae*, *Fusarium oxysporum* f. sp. *niveum*, *Colletotrichum orbiculare*.

Origine Chili (permits)

(d1) The country of origin is free from *Acidovorax avenae* pv *citrulli* (4-NL vrij), *Colletotrichum orbiculare* (4-NL niet vrij), cucumber green mottle mosaic virus (4-NL niet vrij).

EN

(b1) The seeds originate from plants that were inspected during active growing season and found free from *Didymella bryoniae* (6), *Fusarium oxysporum* f. sp. *niveum* (6), squash mosaic virus (6).

OF

(a1) The seeds have been tested in an official laboratory and found free from *Didymella bryoniae* (2), *Fusarium oxysporum* f. sp. *niveum* (2), squash mosaic virus (2).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Didymella bryoniae*, *Fusarium oxysporum* f. sp. *niveum*, squash mosaic virus.

Origine Peru (permits)

(d1) The country of origin is free from *Acidovorax avenae* pv *citrulli* (4-NL vrij), *Fusarium oxysporum* f. sp. *niveum* (4-NL vrij), *Colletotrichum orbiculare* (4-NL niet vrij), cucumber green mottle mosaic virus (4-NL niet vrij), squash mosaic virus (4-NL niet vrij).

EN

(b1) The seeds originate from plants that were inspected during active growing season and found free from *Didymella bryoniae* (6).

OF

(a1) The seeds have been tested in an official laboratory and found free from *Didymella bryoniae* (2).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Didymella bryoniae*.

Coriandrum sp. *

Origines Australië, België, Canada, Denemarken, Duitsland, Finland, Frankrijk, Ierland, Israël, Italië, Japan, Nederland, Nieuw-Zeeland, Oostenrijk, Portugal, Spanje, Taiwan, Verenigd Koninkrijk, Verenigde Staten, Zweden, Zwitserland

Landenoverzicht exporteisen Zaaizaden.

Land: **Israël**

(a1) The seeds have been tested in an official laboratory and found free from *Erwinia carotovora* subsp. *betavasculorum* (2), *Xanthomonas hortorum* pv. *carotae* (2).

OF

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Erwinia carotovora* subsp. *betavasculorum* (6), *Xanthomonas hortorum* pv. *carotae* (6).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Erwinia carotovora* subsp. *betavasculorum*, *Xanthomonas hortorum* pv. *carotae*.

Cosmos sp.

Seeds have undergone an appropriate chemical treatment as specified in the treatment manual. Details vermelden op certificaat in de daartoe bestemde ruimten (12 t/m 17).

Cycas sp.

Seeds have undergone a Phosphine vapour treatment as specified in the treatment manual. Details vermelden op certificaat in de daartoe bestemde ruimten (12 t/m 17).

Seeds are free from pulp.

Cucumis melo *

Origines Australië, België, Canada, Chili, Denemarken, Duitsland, Finland, Frankrijk, Ierland, Israël, Italië, Japan, Nederland, Nieuw-Zeeland, Oostenrijk, Peru, Portugal, Spanje, Taiwan, Turkije, Verenigd Koninkrijk, Verenigde Staten, Zweden, Zwitserland

(c1) The seeds are from an area known to be free from *Acidovorax avenae* subsp. *citrulli* (4-NL vrij; 5)

OF

(a1) The seeds have been tested in an official laboratory and found free from *Acidovorax avenae* subsp. *citrulli* (2),(sample of 10.000 seeds).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Acidovorax avenae* subsp. *citrulli* (sample of 10.000 seeds).

EN

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Fusarium oxysporum* f. sp. *melonis* (6), *Colletotrichum orbiculare* (6), melon necrotic spot virus (6), melon rugose mosaic virus (6), squash mosaic virus (6), tobacco ringspot virus (6).

OF

(a1) The seeds have been tested in an official laboratory and found free from *Fusarium oxysporum* f. sp. *melonis* (2), *Colletotrichum orbiculare* (2), melon necrotic spot virus (2), melon rugose mosaic virus (2), squash mosaic virus (2), tobacco ringspot virus (2).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Fusarium oxysporum* f. sp. *melonis*, *Colletotrichum orbiculare*, melon necrotic spot virus, melon rugose mosaic virus, squash mosaic virus, tobacco ringspot virus.

Cucumis sativus * + Letland, Peru

Origines Australië, België, Canada, Chili, Denemarken, Duitsland, Finland, Frankrijk, Ierland, Israël, Italië, Japan, Nederland, Nieuw-Zeeland, Oostenrijk, Portugal, Spanje, Taiwan, Turkije, Verenigd Koninkrijk, Verenigde Staten, Zweden, Zwitserland

(c1) The seeds are from an area known to be free from *Acidovorax avenae* subsp. *citrulli* (4-NL vrij; 5).

OF

(a1) The seeds have been tested in an official laboratory and found free from *Acidovorax avenae* subsp. *citrulli* (2).

OF

Landenoverzicht exporteisen Zaaizaden.

Land: **Israël**

(a2) The seeds will be tested on-arrival to proof freedom of The seeds have been tested in an official laboratory and found free from *Acidovorax avenae* subsp. *citrulli*.

EN

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Fusarium oxysporum* f. sp. *cucumerinum* (6), *Colletotrichum orbiculare* (6), *Pseudomonas syringae* (6), *Xanthomonas cucurbitae* (6), cucumber green mottle mosaic virus (6) and cucumber leaf spot virus (6).

OF

(a1) The seeds have been tested in an official laboratory and found free from *Fusarium oxysporum* f. sp. *cucumerinum* (2), *Colletotrichum orbiculare* (2), *Pseudomonas syringae* (2), *Xanthomonas cucurbitae* (2), cucumber green mottle mosaic virus (2) and cucumber leaf spot virus (2).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Fusarium oxysporum* f. sp. *cucumerinum*, *Colletotrichum orbiculare*, *Pseudomonas syringae*, *Xanthomonas cucurbitae*, cucumber green mottle mosaic virus and cucumber leaf spot virus.

Origines Peru

(d) The country of origin is free from *Acidovorax avenae* subsp. *citrulli* (4-NL vrij), cucumber green mottle mosaic virus (4-NL niet vrij), *Pseudomonas syringae* (4-NL niet vrij), *Xanthomonas cucurbitae* (4 NL vrij).

OF

(a1) The seeds have been tested in an official laboratory and found free from *Acidovorax avenae* subsp. *citrulli* (2), cucumber green mottle mosaic virus (2), *Pseudomonas syringae* (2), *Xanthomonas cucurbitae* (2).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Acidovorax avenae* subsp. *citrulli*, cucumber green mottle mosaic virus, *Pseudomonas syringae*, *Xanthomonas cucurbitae*.

EN

(b1) The seeds originate from plants that were inspected during the active growing season and found free from cucumber leaf spot virus (6), *Fusarium oxysporum* f. sp. *cucumerinum* (6), *Colletotrichum orbiculare* (6).

OF

(a1) The seeds have been tested in an official laboratory and found free from cucumber leaf spot virus (2), *Fusarium oxysporum* f. sp. *cucumerinum* (2), *Colletotrichum orbiculare* (2).

OF

(a2) The seeds will be tested on-arrival to proof freedom of cucumber leaf spot virus, *Fusarium oxysporum* f. sp. *cucumerinum*, *Colletotrichum orbiculare*.

Origine Letland (permits)

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Colletotrichum orbiculare* (6), cucumber green mottle mosaic virus (6), cucumber leaf spot virus (6), *Fusarium oxysporum* f. sp. *cucumerinum* (6), *Pseudomonas syringae* (6), *Xanthomonas cucurbitae* (6).

OF

(a1) The seeds have been tested in an official laboratory and found free from *Colletotrichum orbiculare* (2), cucumber green mottle mosaic virus (2), cucumber leaf spot virus (2), *Fusarium oxysporum* f. sp. *cucumerinum* (2), *Pseudomonas syringae* (2), *Xanthomonas cucurbitae* (2).

EN

Landenoverzicht exporteisen Zaaizaden.

Land: **Israël**

(c1) The seeds are from an area known to be free from *Acidovorax avenae* subsp. *citrulli* (4-NL vrij; 5).

OF

(a1) The seeds have been tested in an official laboratory and found free from *Acidovorax avenae* subsp. *citrulli* (2).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Acidovorax avenae* subsp. *Citrulli*.

Cucurbita sp. *

Origines Australië, België, Canada, Denemarken, Duitsland, Finland, Frankrijk, Ierland, Israël, Italië, Japan, Nederland, Nieuw-Zeeland, Oostenrijk, Portugal, Spanje, Taiwan, Verenigd Koninkrijk, Verenigde Staten, Zweden, Zwitserland

(c1) The seeds are from an area known to be free from *Acidovorax avenae* subsp. *citrulli* (4-NL vrij; 5).

OF

(a1) The seeds have been tested in an official laboratory and found free from *Acidovorax avenae* subsp. *citrulli* (2).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Acidovorax avenae* subsp. *citrulli*.

EN

(b1) The seeds originate from plants that were inspected during active growing season and found free from *Fusarium oxysporum* f. sp. *melonis* (6), *Colletotrichum orbiculare* (6).

OF

(a1) The seeds have been tested in an official laboratory and found free from *Fusarium oxysporum* f. sp. *melonis* (2), *Colletotrichum orbiculare* (2).

OF.

(a2) The seeds will be tested on-arrival to proof freedom of *Fusarium oxysporum* f. sp. *melonis*, *Colletotrichum orbiculare*.

Cucurbita pepo

Origines Chili (permits)

(c1) The seeds are from an area known to be free from *Acidovorax avenae* subsp. *citrulli* (4-NL vrij; 5), *Colletotrichum orbiculare* (4-NL niet vrij; 5), *Xanthomonas campestris* pv. *cucurbitae* (4-NL vrij; 5), cucumber green mottle mosaic virus (4-NL niet vrij; 5).

EN

(b1) The seeds originate from plants that were inspected during active growing season and found free from *Fusarium solani* (IV), *Phytophthora capsici* (6), squash mosaic virus (6).

OF

(a1) The seeds have been tested in an official laboratory and found free from *Fusarium solani* (I), *Phytophthora capsici* (2), squash mosaic virus (2).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Fusarium solani*, *Phytophthora capsici*, squash mosaic virus.

Cyclamen sp.

(b1) The seeds originate from plants that were inspected during active growing season and found free from *Fusarium oxysporum* f. sp. *cyclaminis* (6).

OF

Landenoverzicht exporteisen Zaaizaden.

Land: **Israël**

(a1) The seeds have been tested in an official laboratory and found free from *Fusarium oxysporum* f. sp. *cyclaminis* (2).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Fusarium oxysporum* f. sp. *cyclaminis*.

Dahlia sp.

(a1) The seeds have been tested in an official laboratory and found free from *Aphelenchoides ritzemabosi* (I), *Gibberella avenaceae* (I).

OF

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Aphelenchoides ritzemabosi* (IV), *Gibberella avenaceae* (IV).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Aphelenchoides ritzemabosi*, *Gibberella avenaceae*.

Daucus carota *

Origines Australië, België, Canada, Denemarken, Duitsland, Finland, Frankrijk, Ierland, Israël, Italië, Japan, Nederland, Nieuw-Zeeland, Oostenrijk, Portugal, Spanje, Taiwan, Verenigd Koninkrijk, Verenigde Staten, Zweden, Zwitserland

The seeds are free from noxious weed seeds of *Cirsium arvense* (I)

Delphinium sp.

(a1) The seeds have been tested in an official laboratory and found free from *Erwinia carotovora* subsp. *atroseptica* (2), *Pseudomonas syringae* pv. *delphinii* (I).

OF

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Erwinia carotovora* subsp. *atroseptica* (6), *Pseudomonas syringae* pv. *delphinii* (I).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Erwinia carotovora* subsp. *atroseptica*, *Pseudomonas syringae* pv. *delphinii*.

Foeniculum sp. *

Origines Australië, België, Canada, Denemarken, Duitsland, Finland, Frankrijk, Ierland, Israël, Italië, Japan, Nederland, Nieuw-Zeeland, Oostenrijk, Portugal, Spanje, Taiwan, Verenigd Koninkrijk, Verenigde Staten, Zweden, Zwitserland

(a1) The seeds have been tested in an official laboratory and found free from *Phomopsis phoenicicola* (I of IV), *Phymatotrichopsis omnivora* (I of IV).

OF

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Phomopsis phoenicicola* (I of IV), *Phymatotrichopsis omnivora* (I of IV).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Phomopsis phoenicicola*, *Phymatotrichopsis omnivora*.

Gaillardia sp.

(a1) The seeds have been tested in an official laboratory and found free from *Alternaria zinniae* (~), *Drechslera* spp. (~), *Pseudocochliobolus pallescens* (~), *Puccinia helianthellae* (~), *Septoria gaillardiae* (~).

OF

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Alternaria zinniae* (~), *Drechslera* spp. (~), *Pseudocochliobolus pallescens* (~), *Puccinia helianthellae* (~), *Septoria gaillardiae* (~).

OF

Landenoverzicht exporteisen Zaaizaden.

Land: **Israël**

(a2) The seeds will be tested on-arrival to proof freedom of *Alternaria zinnia*, *Drechslera* spp., *Pseudocochliobolus pallescens*, *Puccinia helianthellae*, *Septoria gaillardiae*.

Helianthus annuus *

Origines Australië, België, Canada, Chili, Denemarken, Duitsland, Finland, Frankrijk, Ierland, Israël, Italië, Japan, Nederland, Nieuw-Zeeland, Oostenrijk, Portugal, Spanje, Taiwan, Verenigd Koninkrijk, Verenigde Staten, Zweden, Zwitserland

(d2) The seeds originate from plants grown in a country known to be free from *Alternaria helianthi* (4-NL vrij), *Alternaria zinniae* (4-NL niet vrij), *Diaphorthe helianthi* (4-NL vrij), *Leptosphaeria linguistii* (IV), *Septoria helianthi* (4-NL vrij), *Verticillium albo-atrum* (4-NL niet vrij), *Pseudomonas syringae* pv. *tagetis* (4-NL vrij), Sunflower mosaic virus (4-NL vrij), *Orobanche* spp. (IV)

OF

(a1) The seeds have been tested in an official laboratory and found free from *Alternaria helianthi* (2), *Alternaria zinniae* (2), *Diaphorthe helianthi* (2), *Leptosphaeria linguistii* (I), *Septoria helianthi* (2), *Verticillium albo-atrum* (2), *Pseudomonas syringae* pv. *tagetis* (2), Sunflower mosaic virus (2), *Orobanche* spp. (I).

OF

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Alternaria helianthi* (6), *Alternaria zinniae* (6), *Diaphorthe helianthi* (6), *Leptosphaeria linguistii* (IV), *Septoria helianthi* (6), *Verticillium albo-atrum* (6), *Pseudomonas syringae* pv. *tagetis* (6), Sunflower mosaic virus (6), *Orobanche* spp. (IV)

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Alternaria helianthi*, *Alternaria zinniae*, *Diaphorthe helianthi*, *Leptosphaeria linguistii*, *Septoria helianthi*, *Verticillium albo-atrum*, *Pseudomonas syringae* pv. *tagetis*, Sunflower mosaic virus, *Orobanche* spp

AND

Seeds have undergone a Phosphine vapour treatment as specified in the treatment manual. *Let op: behandeling vermelden op fytcertificaat in de daartoe bestemde ruimten (12 t/m 17).*

Lactuca sativa *

Origines Australië, België, Canada, Denemarken, Duitsland, Finland, Frankrijk, Ierland, Israël, Italië, Japan, Nederland, Nieuw-Zeeland, Oostenrijk, Portugal, Spanje, Taiwan, Verenigd Koninkrijk, Verenigde Staten, Zweden, Zwitserland

(a1) The seeds have been tested in an official laboratory and found free from *Verticillium albo-atrum* (I), *Xanthomonas campestris* pv. *vitians* (2).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Verticillium albo-atrum*, *Xanthomonas campestris* pv. *vitians*.

EN

(a1) The seeds have been tested in an official laboratory (seed sample of 10.000) and found free from lettuce mosaic virus (2). (*Let op: Israël eist een monster van 10.000 zaden voor de toets op lettuce mosaic virus.*)

OF

(a2) The seeds will be tested on-arrival to proof freedom of lettuce mosaic virus.

Lathyrus odoratus

~~(b2 of a1 of a2) + *Ascochyta lathyri* (IV), *Rhodococcus fascians* (2; 6), pea mosaic virus = bean yellow mosaic virus (IV of I).~~

Landenoverzicht exporteisen Zaaizaden.

Land: **Israël**

(a1) The seeds have been tested in an official laboratory and found free from *Ascochyta lathyri* (IV), *Rhodococcus fascians* (2) pea mosaic virus = bean yellow mosaic virus (~~2IV of I~~).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Ascochyta lathyri* (~~IV~~), *Rhodococcus fascians* (~~2; 6~~), pea mosaic virus = bean yellow mosaic virus (~~IV of I~~).

OF

(b2) The seeds originate from a field inspected during the active growing season and found free from *Ascochyta lathyri* (IV), *Rhodococcus fascians* (~~2; 6~~), pea mosaic virus = bean yellow mosaic virus (~~6IV of I~~).

Petroselinum sp. *

Origines Australië, België, Canada, Denemarken, Duitsland, Finland, Frankrijk, Ierland, Israël, Italië, Japan, Nederland, Nieuw-Zeeland, Oostenrijk, Portugal, Spanje, Taiwan, Verenigd Koninkrijk, Verenigde Staten, Zweden, Zwitserland

(a1) The seeds have been tested in an official laboratory and found free from *Phymatotrichopsis omnivora* (IV of I), *Phytophthora cryptogea* (2), *Pythium debaryanum* (IV of I), strawberry latent ringspot virus (2).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Phymatotrichopsis omnivora*, *Phytophthora cryptogea*, *Pythium debaryanum*, strawberry latent ringspot virus.

OF

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Phymatotrichopsis omnivora* (IV of I), *Phytophthora cryptogea* (6), *Pythium debaryanum* (IV of I), strawberry latent ringspot virus (6).

EN

The seeds are free from noxious weed seeds of *Cirsium arvense* (I).

Petunia sp.

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Erwinia chrysanthemi* (6), impatiens necrotic spot virus (6), potato spindle tuber viroid (6), tobacco ringspot virus (6), tomato aspermy virus (6), tomato black ring virus (6), tomato bushy stunt virus (6), tomato ringspot virus (6).

OF

(a1) The seeds have been tested in an official laboratory and found free from *Erwinia chrysanthemi* (2), impatiens necrotic spot virus (2), potato spindle tuber viroid (2), tobacco ringspot virus (2), tomato aspermy virus (2), tomato black ring virus (2), tomato bushy stunt virus (2), tomato ringspot virus (2).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Erwinia chrysanthemi*, impatiens necrotic spot virus, potato spindle tuber viroid, tobacco ringspot virus, tomato aspermy virus, tomato black ring virus, tomato bushy stunt virus, tomato ringspot virus.

Phaseolus sp. *

Origines Australië, België, Canada, Denemarken, Duitsland, Finland, Frankrijk, Ierland, Israël, Italië, Japan, Nederland, Nieuw-Zeeland, Oostenrijk, Portugal, Spanje, Taiwan, Verenigd Koninkrijk, Verenigde Staten, Zweden, Zwitserland

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Ditylenchus dipsaci* (IV), *Colletotrichum lindemuthianum* (6), *Colletotrichum truncatum* (6), *Diaporthe phaseolorum* (IV), *Phoma exigua* (6), *Phyllosticta phaseolina* (IV), *Curtobacterium flaccumfaciens* pv. *flaccumfaciens* (6), *Pseudomonas syringae* pv. *phaseolicola* (6), *Xanthomonas axonopodis* pv. *phaseoli* (6), bean common mosaic virus (6), tobacco streak virus (6).

OF

Landenoverzicht exporteisen Zaaizaden.

Land: **Israël**

(a1) The seeds have been tested in an official laboratory and found free from *Ditylenchus dipsaci* (I), *Colletotrichum lindemuthianum* (2), *Colletotrichum truncatum* (2), *Diaporthe phaseolorum* (I), *Phoma exigua* (2), *Phyllosticta phaseolina* (I), *Curtobacterium flaccumfaciens* pv. *flaccumfaciens* (2), *Pseudomonas syringae* pv. *phaseolicola* (2), *Xanthomonas axonopodis* pv. *phaseoli* (2), bean common mosaic virus (2), tobacco streak virus (2).

OF

(a2) The seeds will be tested on-arrival to proof freedom of from *Ditylenchus dipsaci*, *Colletotrichum lindemuthianum*, *Colletotrichum truncatum*, *Diaporthe phaseolorum*, *Phoma exigua*, *Phyllosticta phaseolina*, *Curtobacterium flaccumfaciens* pv. *flaccumfaciens*, *Pseudomonas syringae* pv. *phaseolicola*, *Xanthomonas axonopodis* pv. *phaseoli*, bean common mosaic virus, tobacco streak virus.

Pisum sp. *

Origines Australië, België, Canada, Denemarken, Duitsland, Finland, Frankrijk, Ierland, Israël, Italië, Japan, Nederland, Nieuw-Zeeland, Oostenrijk, Portugal, Spanje, Taiwan, Verenigd Koninkrijk, Verenigde Staten, Zweden, Zwitserland

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Ascochyta pinodes* (6), pea enation mosaic virus (6), pea seed-borne mosaic virus (IV)

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Ascochyta pinodes*, pea enation mosaic virus, pea seed-borne mosaic virus.

EN

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Pseudomonas syringae* pv. *pisi* (I + IV).

EN

(a1) The seeds have been tested in an official laboratory and found free from *Pseudomonas syringae* pv. *pisi* (I + IV).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Pseudomonas syringae* pv. *pisi*.

EN

The seeds have undergone a Phosphine vapour treatment and an appropriate chemical treatment as specified in the treatment manual.

Let op: behandeling vermelden op het FC.

Primula sp.

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Phyllosticta primulicola* (IV), *Pseudomonas syringae* pv. *primulae* (6).

OF

(a1) The seeds have been tested in an official laboratory and found free from *Phyllosticta primulicola* (I), *Pseudomonas syringae* pv. *primulae* (2).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Phyllosticta primulicola*, *Pseudomonas syringae* pv. *primulae*.

Ranunculus asiaticus

(b1 of a1 of a2) + *Colletotrichum acutatum* (IV of I).

Raphanus sativus

Landenoverzicht exporteisen Zaaizaden.

Land: **Israël**

Origines Australië, België, Canada, Denemarken, Duitsland, Finland, Frankrijk, Ierland, Israël, Italië, Japan, Nederland, Nieuw-Zeeland, Oostenrijk, Portugal, Spanje, Taiwan, Verenigd Koninkrijk, Verenigde Staten, Zweden, Zwitserland

Geen bijschrijvingen.

Overige origines (permits)

(a1) The seeds have been tested in an official laboratory and found free from *Sclerotinia* spp. (2; 21)

(a2) The seeds will be tested on-arrival to proof freedom of *Sclerotinia* spp.

Salpiglossis sp. *

Origines Australië, België, Canada, Denemarken, Duitsland, Finland, Frankrijk, Ierland, Israël, Italië, Japan, Nederland, Nieuw-Zeeland, Oostenrijk, Portugal, Spanje, Taiwan, Verenigd Koninkrijk, Verenigde Staten, Zweden, Zwitserland

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Clavibacter michiganensis* subsp. *michiganensis* (6).

OF

(a1) The seeds have been tested in an official laboratory and found free from *Clavibacter michiganensis* subsp. *michiganensis* (2).

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Clavibacter michiganensis* subsp. *Michiganensis*.

Solanum lycopersicum

Alle origines (m.u.v. Kenia, Peru, Zuid-Afrika) (permits)

Zie ook 'Certificeringseisen'.

Eis zonder bijschrijving:

Partij vrij van tomato brown rugose fruit virus (2).

EN

The seeds are free from *Fusarium oxysporum* f. sp. *lycopersici* race III (2; 4 NL vrij; 5; 6; 9), *Clavibacter michiganensis* (2; 4-NL vrij; 5; 6; 9), *Erwinia carotovora* subsp. *carotovora* (6), *Pseudomonas corrugata* (2; 4-NL niet vrij; 5; 6; 9; 13), *Pseudomonas syringae* pv. *tomato* (2; 4 NL vrij; 5; 6; 9), *Xanthomonas vesicatoria* (2; 4 NL vrij; 5; 6; 9), *pepino mosaic virus* (2; 4 NL niet vrij; 5; 6; 9; 13), *potato spindle tuber viroid* (2; 4 NL niet vrij; 5; 6; 9), *tobacco mosaic virus* (2; 4 NL niet vrij; 5; 6; 9; 13), *tomato black ring virus* (2; 4-NL vrij; 5; 6; 9; 13), *tobacco mild green mosaic virus* (IV), *tomato black ring virus* (98), *tomato bushy stunt virus* (98), *tomato ringspot virus* (98).

OF

(a1) The seeds have been tested in an official laboratory and found free from *Fusarium oxysporum* f. sp. *lycopersici* race III (2), *Clavibacter michiganensis* (2), *Erwinia carotovora* subsp. *carotovora* (2), *Pseudomonas corrugata* (2), *Pseudomonas syringae* pv. *tomato* (2), *Xanthomonas vesicatoria* (2), *pepino mosaic virus* (2), *potato spindle tuber viroid* (2), *tobacco mosaic virus* (2), *tomato mosaic virus* (2), *tobacco mild green mosaic virus* (I), *tomato black ring virus* (2), *tomato bushy stunt virus* (2), *tomato ringspot virus* (2).

EN

The seeds have been acid extracted.

Origines Kenia, Peru, Zuid-Afrika (permits)

Geen bijschrijvingen.

Solanum melongena

Allee origines (m.u.v. China).

Zie ook bij certificeringseisen.

Landenoverzicht exporteisen Zaaizaden.

Land: **Israël**

The production area is free from *Fusarium oxysporum* f. sp. *melongenae* (4-NL niet vrij; 5), *Phomopsis vexans* (4-NL vrij; 5), *Verticillium albo-atrum* (4-NL niet vrij; 5), *Ralstonia solanacearum* (4-NL niet vrij; 5), eggplant mosaic virus (4-NL vrij; 5), Tobacco ringspot virus (4-NL niet vrij; 5).

OF

The consignment was checked during active growth and it was found free from *Fusarium oxysporum* f. sp. *melongenae* (6), *Phomopsis vexans* (6), *Verticillium albo-atrum* (6), *Ralstonia solanacearum* (6), eggplant mosaic virus (6), Tobacco ringspot virus (6).

OF

Based on laboratory tests the consignment was found free from *Fusarium oxysporum* f. sp. *melongenae* (2), *Phomopsis vexans* (2), *Verticillium albo-atrum* (2), *Ralstonia solanacearum* (2), eggplant mosaic virus (2), Tobacco ringspot virus (2). (*Let op: labuitslag meesturen.*)

The seeds have undergone an appropriate chemical treatment as specified in the treatment manual. *Let op: behandeling vermelden op het fytocertificaat in de daartoe bestemde ruimten (12 t/m 17).*

Origine China (permits)

Geen bijschrijvingen.

Solanum tuberosum

Origine Nederland (permits)

Based on official sampling and testing, the following organism does not occur in the cultivation chain of potato in the Netherlands: potato spindle tuber viroid (2; 9).

Let op: deze bijschrijving wijkt af van de door Israel gevraagde eis: The country of production is free from potato spindle tuber viroid.

EN

The place of production and seed crop were inspected during the active growth and found free from: *Synchytrium endobioticum* (6), *Clavibacter sepedonicus* (6), *Ralstonia solanacearum* (6).

Tagetes sp.

(a1) The seeds have been tested in an official laboratory and found free from *Fusarium oxysporum* f. sp. *callistephi* (2), *Pseudomonas syringae* pv. *tagetis* (2).

OF

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Fusarium oxysporum* f. sp. *callistephi* (6), *Pseudomonas syringae* pv. *tagetis* (6)

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Fusarium oxysporum* f. sp. *callistephi*, *Pseudomonas syringae* pv. *tagetis*.

Tropaeolum majus

(a1) The seeds have been tested in an official laboratory and found free from *Rhodococcus fascians* (2).

OF

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Rhodococcus fascians* (6).

OF

The seeds will be tested on-arrival to proof freedom of *Rhodococcus fascians*.

EN

The seeds have undergone a hot water treatment as specified in the treatment manual. *Let op: behandeling vermelden op het FC.*

Landenoverzicht exporteisen Zaaizaden.

Land: **Israël**

Zea mays

Origine Verenigde Staten (permits)

(d1) The country of origin is free from *Sclerophthora macrospora* (4-NL vrij), *Colletotrichum graminicola* (4-NL vrij), *Stenocarpella maydis* (4-NL vrij).

OF

(d3) The place of production was officially inspected during the growing season and found free from *Sclerophthora macrospora* (6), *Colletotrichum graminicola* (6), *Stenocarpella maydis* (6).

Zinnia sp.

(a1) The seeds have been tested in an official laboratory and found free from *Alternaria zinniae* (2), *Xanthomonas campestris* pv. *zinniae* (2).

OF

(b1) The seeds originate from plants that were inspected during the active growing season and found free from *Alternaria zinniae* (6), *Xanthomonas campestris* pv. *zinniae* (6)

OF

(a2) The seeds will be tested on-arrival to proof freedom of *Alternaria zinniae*, *Xanthomonas campestris* pv. *zinniae*.

Hoewel de NVWA dit document op zorgvuldige wijze en naar beste weten heeft samengesteld, kan niet worden ingestaan voor de juistheid en volledigheid van de beschikbaar gestelde informatie. Aan de beschikbaar gestelde informatie kunnen geen rechten worden ontleend. Een afdruk kan verouderd zijn. Een actuele versie is op de website van NVWA beschikbaar.